

## **PARKINSON'S DISEASES**

[Theta-phase closed-loop stimulation induces motor paradoxical responses in the rat model of Parkinson disease.](#)

Cordon I, Nicolás MJ, Arrieta S, Alegre M, Artieda J, Valencia M. Brain Stimul. 2018 Jan - Feb;11(1):231-238. doi: 10.1016/j.brs.2017.10.004.

[GPR55: A therapeutic target for Parkinson's disease?](#)

Celorio M, Rojo-Bustamante E, Fernández-Suárez D, Sáez E, Estella-Hermoso de Mendoza A, Müller CE, Ramírez MJ, Oyarzábal J, Franco R, Aymerich MS. Neuropharmacology. 2017 Oct;125:319-332. doi: 10.1016/j.neuropharm.2017.08.017.

[Fatty acid amide hydrolase inhibition for the symptomatic relief of Parkinson's disease.](#)

Celorio M, Fernández-Suárez D, Rojo-Bustamante E, Echeverry-Alzate V, Ramírez MJ, Hillard CJ, López-Moreno JA, Maldonado R, Oyarzábal J, Franco R, Aymerich MS. Brain Behav Immun. 2016 Oct;57:94-105. doi: 10.1016/j.bbi.2016.06.010.

[Monoacylglycerol lipase inhibitor JZL 184 is neuroprotective and alters glial cell phenotype in the chronic MPTP mouse model.](#)

Fernández-Suárez D, Celorio M, Riezu-Boj JI, Ugarte A, Pacheco R, González H, Oyarzábal J, Hillard CJ, Franco R, Aymerich MS. Neurobiol Aging. 2014 Nov;35(11):2603-2616. doi: 10.1016/j.neurobiolaging.2014.05.021.

[Abnormal functional connectivity between motor cortex and pedunculopontine nucleus following chronic dopamine depletion.](#)

Valencia M, Chavez M, Artieda J, Bolam JP, Mena-Segovia J. J Neurophysiol. 2014 Jan;111(2):434-40. doi: 10.1152/jn.00555.2013.

[Cortical oscillations scan using chirp-evoked potentials in 6-hydroxydopamine rat model of Parkinson's disease.](#)

Pérez-Alcázar M, Nicolás MJ, Valencia M, Alegre M, López-Azcárate J, Iriarte J, Artieda J. Brain Res. 2010 Jan 15;1310:58-67. doi: 10.1016/j.brainres.2009.11.031.